

## Should local treatment of breast ductal carcinoma *in situ* be the same as the treatment of early invasive breast cancer

Sylwia Grodecka-Gazdecka

Heterogenicity of breast ductal carcinoma *in situ* gives rise to opposing proposals concerning its treatment — ranging from attempts to recommend the *watch and wait* strategy in low risk forms ending with the currently binding standards of treatment of DCIS in the way identical as early invasive cancer in the high risk. Arguments for the treatment of ductal carcinoma *in situ* in the same way as patients with early invasive cancer have been presented. These arguments comprise: unknown natural history of untreated DCIS, high risk of undervaluation of the invasive component in the core-needle biopsy, the increase of recurrence risk with the progress of time, lack of verified separators of the groups with the risk of adverse course of the disease, the results of the clinical studies confirming the justification of combined local treatment and the proof that the clinical course of DCIS is the same as early invasive breast cancer, and, first and foremost, the fact that there are no clinical studies which could justify a limitation of the treatment scope.

NOWOTWORY J Oncol 2017; 67, 1: 74–78

**Key words:** DCIS — breast ductal carcinoma *in situ*, cancer heterogenicity, local treatment, watch and wait strategy, clinical studies

Breast ductal carcinoma *in situ* (DCIS) is characterised with a malignant transformation of the epithelium of the ducts, which is limited to the epithelia and myoepithelial layers without crossing the basal cell membrane. This type of cancer is regarded as a transitional stage between the normal breast tissue and invasive cancer. It is characterised with a significant diversity of the morphological, immunochemical, molecular and clinical picture [1]. Depending on the tissue architecture, DCIS can be defined as solid, cribriform, papillary and micropapillary; whilst depending on the malignancy grade, as: poorly-, moderately- and highly differentiated and — depending on the presence of the *comedo* type necrosis — as *comedo* type carcinoma (with a more aggressive clinical course) and non-*comedo* type carcinoma. Intraductal spread of the disease in connection with an irregular routing of the ducts and difficulties in macroscopic evaluation of the scope of the lesions illustrates well the deceptive character of the disease and the widely understood heterogenicity of the DCIS requires some si-

gnificant evaluation of the therapeutic management [2]. The risk of development of an invasive form of cancer, which — depending on the subtype of DCIS — is 20–30% within 10 years and is 15 times greater than the average risk of breast cancer morbidity in the general population [3] of key importance for the choice of the scope of treatment.

The scale of the problem can be illustrated with the fact that since the 1970s there has been a continued increase in diagnoses of ductal carcinoma *in situ* due to the popularisation of screening examinations and more advanced diagnostic techniques. Currently DCIS makes up about 20–25% of breast cancers in the Western countries and in Poland 7–10% of new diagnoses per year [4]. In spite of the increasing number of DCIS diagnoses and early invasive cancers, the number of diagnoses of advanced breast cancer have not dropped, which remains a paradox [5].

Diagnosis of ductal breast carcinoma *in situ* is made in 90% of cases during a screening mammography, whilst, in a clinical examination such diagnosis is made much more

rarely, usually only when a tumour in the breast or an exudation from a papilla is found. DCIS comprises a vast range of symptoms, from some slight low-risk lesions to eventually the involvement of extensive breast areas with lesions of high malignancy potential. The most frequent symptom of the disease consists in lesions, visible in mammography as accumulations of malignant micro-calcifications, which can be uni- or bilateral, frequently multifocal. The MRI mammography, useful in particular for the evaluation of the multifocality of neoplastic lesions and a more accurate estimation of the tumour size, is more sensitive than classical mammography. A microscopic diagnosis is usually made from the specimen collected in a mammotomy biopsy or — less frequently — in the surgical specimen, in which DCIS may occur independently or co-exist with an invasive carcinoma.

The primary objective in the DCIS treatment is to reduce the number of recurrences and to decrease the number of recurrences with an invasive component. The determination of the groups with the risk of an adverse course of the disease is of essential importance for the choice of optimum therapeutic management. The DCIS heterogeneity thus gives rise to opposing proposals concerning its treatment — from attempts to recommend the watch and wait strategy in low risk forms to currently binding standards of treatment of DCIS in the same way as early invasive cancer in the high risk group [6].

The scope of treatment of DCIS ranges from the resection of the neoplastic lesion, through an excision with an adjuvant hormonotherapy, excision with the sentinel node biopsy and adjuvant radiotherapy, excision with the sentinel node biopsy and adjuvant radiotherapy with hormonotherapy, simple mastectomy, simple mastectomy with the sentinel node biopsy, subcutaneous mastectomy with the sentinel node biopsy and immediate reconstruction to end with the subcutaneous mastectomy with the sentinel node biopsy and immediate reconstruction and adjuvant hormonotherapy. The choice of the optimum scope of loco-regional treatment is usually assisted with the assessment of the prognostic value of the Van Nuys Prognostic Index (VNPI), worked out in 2002 or VNPI/SCI, corrected in 2009 [7]. Amongst the tools recommended for a more specific prediction of the course of the disease, are also genetic tests, including Oncotype DX and the examination of the *BRCA1* and *BRCA2* gene mutation [8, 9].

The most popular locoregional management is organ sparing treatment, consisting of the resection of the neoplastic lesion with an adjuvant radiotherapy. The indications for the sentinel node biopsy are limited to patients with a high risk of the presence of a malignant component, (large areas of malignant calcifications, location in the tail of Spence, planned mastectomy) [10]. In spite of the predominance of breast sparing treatment, the published data points to

significant differentiation of treatment choices resulting from the attitudes of the patients and the experience of the therapeutic team [11, 12]. In the USA, after the period of a decrease in the number of mastectomies in 1998–2004 (from 36 to 28%), since 2011 a further growth in the number of breast amputations, following the diagnose of DCIS has been observed, correlating additionally with the increase of the contralateral breast amputation. This concerns mostly younger patients, *BRCA1* gene carriers, in particular, in families with a family history burdened with ovarian cancer [9]. An additional argument for this option is the reluctance for revision surgeries necessary to obtain a cancer-free margin, chimeric course of the disease in spite of a lack of any signs of invasion, the possibility to avoid radiotherapy and access to the procedures of immediate breast reconstruction [13–15]. This escalation of the scope of surgical treatment is one of the mechanisms driving the debate on the necessity to look for decision factors, allowing the optimisation of the treatment of breast ductal carcinoma *in situ*.

The basic arguments for the treatment of ductal carcinoma *in situ* in the same way as early invasive cancer comprise:

- unknown natural history of untreated DCIS [16];
- high risk of undervaluation of the invasive component in the core-needle biopsy [10, 16–18];
- increase of recurrence risk with the progress of time [3, 19–21];
- lack of verified separators of the groups with the risk of adverse course of the disease [1, 2, 20];
- the results of the clinical studies confirming the justification of combined local treatment [22–26];
- and the proof that the clinical course of DCIS is the same as early invasive breast cancer [27, 28];
- the lack of clinical studies which could justify a limitation of the treatment scope [28–30].

Given the fact that a large share of ductal carcinoma *in situ* is diagnosed as a small lesion seen only in a mammography image, and then treated with a mammotomy biopsy, a substantial part of DCIS is resected during this procedure. Thus, the natural history of untreated DCIS remains unknown [16].

According to the published data, in 10 to 50% patients with ductal carcinoma *in situ* diagnosed during a core-needle or mammotomy biopsy, there is a risk of the underestimation of the co-existence of an invasive component, which is illustrated by microscopic evaluation performed after the surgical resection of the lesion [2, 10, 17, 18]. In the opinion of the authors of this meta-analysis, which comprised 7350 subject with a diagnosis of DCIS established with a core-needle biopsy, a consequence of underestimation of the risk of the presence of an invasive component was the delay of the correct diagnosis and treatment in one out of five cases. The presence of invasive cancer was found in 1738 patients from the analysed group [18]. In the studies

in which an invasive component was diagnosed in 40% of patients, the following factors were regarded as a risk: the presence of tissue mass in the preoperative MRI mammography, the involvement of the nipple-areolar complex, a large heterogeneity of the lesion and HER2 overexpression [10, 17]. The recurrence risk increases with the progress of time and, given the unknown natural history of untreated DCIS, the observations concerning disease recurrence are a source of knowledge of DCIS biology and the effects on the therapy applied. In one-centre retrospective study comprising 200 patients, within an 8-year observation period, in 25 patients (12.5%) a disease recurrence was diagnosed, and the risk factors were the young age, tumour size and grade [3]. In a prospective multicentre trial, ECOG-ACRIN E5194 with an average observation period of 12.3 years, where the minimum cancer-free margin was more than 3 mm, special attention was paid to the group of patients with a potentially low recurrence risk low-grade DCIS. In this group, comprising 561 patients, also 14.4% of recurrence cases were found (vs 24.6% in the high-grade group), including 7.5% invasive recurrences (vs 13.4% in the high-grade group) [19]. In the group of patients with an almost 20-year observation period, after the treatment of asymptomatic DCIS, found in the mammographic picture, 16.3% of recurrences were found,  $\frac{3}{4}$  of which were invasive and the independent risk factor was an age below 45 years [24].

In spite of the probability of a different course of disease depending on the group with a low or high tumour grade, there are no verified and well documented risk separators. The knowledge of the molecular aspect of DCIS will allow for a better understanding of heterogeneity of this type of cancer and its significance for the clinical picture of a disease. For the time being the studies concentrate on the analysis of epigenetic modifications such as DNA methylation and the changes within miRNA, which play a significant role in genetic disorders. Additionally, the tumour increase and invasion are facilitated by a distorted tumour microenvironment, whose fibroblasts and macrophages excrete growth factors and angiogenesis stimulating factors [1]. Another group of molecular studies concern the relationships of the expression of selected biomarkers and the prognoses, where the triple-positive DCIS (p16, COX-2, Ki67) is significantly related to a higher risk of the occurrence of invasive breast cancer [31].

In the light of the recent studies whose results may in future foster the modification of clinical decisions, it must be concluded that the current state of knowledge, providing strong evidence justifying the local treatment of breast DCIS in the same way as early invasive form of this cancer [22]. With regards to the strength of the arguments from clinical studies, an identical standpoint is presented in the current recommendations of NCCN, AGO, ESMO or recommendations of the Polish Society of Surgical Oncology, currently

prepared for publication [32–34]. The arguments for the positive answer were provided, among others, by the results of the NSABP studies: B-17 and B-24 in their section concerning DCIS. Within more than a 15-year observation period, 490 recurrences were observed, including 263 (53.7%) invasive cases, which were related to a higher risk of death. A significant reduction of recurrences was confirmed in the group of patients undergoing adjuvant radiotherapy (and also among those treated with tamoxifen). The share of contralateral breast cancer, within the period of 15 years was 10% in the studied groups, whilst it was reduced to 7.3% after the addition of tamoxifen to the therapy. It was proven that multiple year prognoses after surgical treatment is very good [23]. The recurrence risk factors were the young age of the patients and the lack of a tumour free margin [23, 35].

A definition of an adequate margin still remains under discussion. What is beyond all doubt, however, is the significance of the cancer-free margin as the strongest risk factor of local recurrence [36–39]. In accordance with the St. Gallen recommendations, a cancer free margin means the lack of cells of invasive or pre-invasive cancer in the excision line (“no ink on invasive tumour or DCIS”) [40]. In the meta-analysis of 20 studies, published in August 2016 and concerning as many as 7883 patients with a mean observation period of 78.3 months, a definite standpoint of three opinion-forming American scientific societies (SSO, ASTRO, ASCO), that a standard in the treatment of DCIS is a margin not smaller than 2 mm in patients undergoing surgical intervention and a radical adjuvant radiotherapy (WBRT) [41].

The application of adjuvant radiotherapy in breast conserving surgery in the patients with DCIS has a proven significance for decreasing the recurrence risk [26]. The prospective clinical study, RTOG 9804, proved the benefit in adjuvant radiotherapy also in the group of patients with good prognoses (low and medium recurrence risk in the patients with negative surgical margin). During the mean observation period of 7.17 years in the group of patients without irradiation, there was 42% cases of disease recurrence, whilst the addition of radiotherapy resulted in a reduction of the recurrence risk from 6.7% to 0.9% [25]. The analysis of data coming from the SEER register (Surveillance, Epidemiology and End Results), concerning the population of 56968 people treated for DCIS with an average observation period of 91 months, confirmed the beneficial effect of adjuvant radiotherapy on the overall survival period throughout the entire study group, yet this effect was expressed in the strongest way in younger patients, without or with a low content of oestrogen receptors in the nuclei of the cancer cells [42].

Strong arguments for the treatment of DCIS in the same way as early invasive cancer are also presented in the publications in which the analysis of the data from the SEER register concerning 108 196 women with a diagnosis of DCIS confirms the fact that the clinical course of the disease in

both groups of patients is similar [27, 28]. One of the most essential contra-arguments against the reduction of local treatment, even in the group defined as *low risk* DCIS, is the lack of results of such randomised clinical studies as LORIS in Great Britain, which is a prospective study comparing surgical treatment (plus adjuvant radiotherapy or hormonotherapy) with an active observation, which was supposed to foster better understanding of the natural history of untreated DCIS [29]. In 2015 also, the LORD study was started (the third phase study EORTC-BCG 1401/BOOG 2014-04), in which in 1240 patients with a diagnosis of low risk DCIS, and with a 10 year follow-up period and active observation (watch and wait strategy) will be compared with treatment considered so far to be the standard [30].

In the light of the presented facts, the authors of this publication wish to sustain their standpoint that a local treatment of a patient with a ductal carcinoma *in situ* should be the same as the treatment of patients with early invasive cancer, as there is strong evidence for the effectiveness of the local treatment of DCIS, whilst the abandonment of surgical treatment and the treatment combined with radiotherapy remains premature without any data from the prospective clinical studies which could confirm the safety of such procedure.

**Conflict of interest:** none declared

**Prof. Sylwia Grodecka-Gazdecka, MD, PhD**

*Oncology Chair and Clinic*

*Poznań University of Medical Sciences*

*Szamarzewskiego 82/84*

*60–569 Poznań, Poland*

*e-mail: sylwia.grodecka-gazdecka@skpp.edu.pl*

*Received & Accepted: 6 Sept 2016*

Based on the presentation at the IV Annual Conference of the *Nowotwory Journal of Oncology*, 'Oncological Debates', held in Warszawa, 8–9th April 2016

## References

- Mardekian SK, Bombonati A, Palazzo JP. Ductal carcinoma in situ of the breast: the importance of morphologic and molecular interactions. *Hum Pathol* 2016; 49: 114–123.
- Pang JM B, Gorringer KL, Fox SB. Ductal carcinoma in situ — update on risk assessment and management. *Histopathology* 2016; 68: 96–109.
- van Mackelenbergh MT, Lindner CM, Heilmann T et al. Impact of histopathological factors, patient history and therapeutic variables on recurrence-free survival after ductal carcinoma in situ: 8-year follow-up and questionnaire survey. *Geburtshilfe Frauenheilkd* 2016; 76: 46–52.
- Didkowska J, Wojciechowska U. Nowotwory piersi w Polsce i Europie — populacyjny punkt widzenia. *Nowotwory J Oncol* 2013; 63: 111–118.
- DeSantis C, Howlader N, Cronin KA et al. Breast cancer incidence rates in U.S. women are no longer declining. *Cancer Epidemiol Biomarkers Prev* 2011; 20: 733–739.
- Kuerer HM. Ductal carcinoma in situ: treatment or active surveillance? *Expert Rev Anticancer Ther* 2015; 15: 777–785.
- Silverstein MJ. The University of Southern California/Van Nuys prognostic index for ductal carcinoma in situ of the breast. *Am J Surg* 2003; 186: 337–343.
- Rakovitch E, Nofech-Mozes S, Hanna W et al. A population-based validation study of the DCIS score predicting recurrence risk in individuals treated by breast conserving therapy alone. *Breast Cancer Res Treat* 2015; 152: 389–398.
- Elsayegh N, Kuerer HM, Lin H et al. Predictors that influence contralateral prophylactic mastectomy election among women with ductal carcinoma in situ who were evaluated for BRCA genotyping testing. *Ann Surg Oncol* 2014; 21: 3466–3472.
- Tunon-de-Lara C, Chauvet MP, Baranzelli MC et al. The role of sentinel lymph node biopsy and factors associated with invasion in extensive DCIS of the breast treated by mastectomy: the Cinnamon Prospective Multicenter Study. *Ann Surg Oncol* 2015; 22: 3853–3860.
- Worni M, Akushevich I, Greenup R et al. Trends in treatment patterns and outcomes for ductal carcinoma in situ. *J Natl Cancer Inst* 2015; 107: djv 263.
- Manuu GS, Bettencourt-Silva JH, Ahmed F et al. A nationwide cross-sectional survey of UK breast surgeons' views on management of ductal carcinoma in situ. *Int J Breast Cancer* 2015; 104231. doi:10.1155/2015/104231.
- Rutter CE, Park HS, Killelea BK et al. Growing use of mastectomy for ductal carcinoma in situ of the breast among young women in the United States. *Ann Surg Oncol* 2015; 22: 2378–2386.
- Ashfaq A, McGhan LJ, Pockaj BA et al. Impact of breast reconstruction on the decision to undergo contralateral prophylactic mastectomy. *Ann Surg Oncol* 2014; 21: 2934–2940.
- Habermann EB, Thomsen KM, Hieken TJ et al. Impact of availability of immediate breast reconstruction on bilateral mastectomy rates for breast cancer across the United States: data from the nationwide inpatient sample. *Ann Surg Oncol* 2014; 21: 3290–3296.
- Virnig BA, Tuttle TM, Shamlivan T et al. Ductal carcinoma in situ of the breast: a systematic review of incidence, treatment, and outcomes. *J Natl Cancer Inst* 2010; 102: 170–178.
- Lee CW, Wu HK, Lai HW et al. Preoperative clinicopathological factors and breast magnetic resonance imaging features can predict ductal carcinoma in situ with invasive components. *Eur J Radiol* 2016; 85: 780–789.
- Brennan ME, Turner RM, Ciatto S et al. Ductal carcinoma in situ at core-needle biopsy: meta-analysis of underestimation and predictors of invasive breast cancer. *Radiology* 2011; 260: 119–128.
- Solin LJ, Gray R, Hughes LL et al. Surgical excision without radiation for ductal carcinoma in situ of the breast: 12-year results from the ECOG-ACRIN E5194 study. *J Clin Oncol* 2015; 33: 3938–3944.
- Dereere E, Papadimitriou K, Tjalma W et al. Ductal carcinoma in situ: a disease entity that merits more recognition. *Minerva Chir* 2015; 70: 231–239.
- Cheung S, Booth ME, Kearins O et al. Risk of subsequent invasive breast cancer after diagnosis of ductal carcinoma in situ (DCIS). *Breast* 2014; 23: 807–811.
- Merrill AL, Esserman L, Morrow M. Clinical decisions. Ductal carcinoma in situ. *N Engl J Med* 2016; 374: 390–392.
- Wapnir IL, Dignam JJ, Fisher B et al. Long-term outcomes of invasive ipsilateral breast tumor recurrences after lumpectomy in NSABP B-17 and B-24 randomized clinical trials for DCIS. *J Natl Cancer Inst* 2011; 103: 478–488.
- Wilkinson JB, Vicini FA, Shah Ch et al. Twenty-year outcomes after breast-conserving surgery and definitive radiotherapy for mammographically detected ductal carcinoma in situ. *Ann Surg Oncol* 2012; 19: 3785–3791.
- McCormick B, Winter K, Hudis C et al. RTOG 9804: A prospective randomized trial for good-risk ductal carcinoma in situ comparing radiotherapy with observation. *J Clin Oncol* 2015; 33: 709–715.
- Early Breast Cancer Trialists', Correa C, McGale P, Taylor C et al. Overview of the randomized trials of radiotherapy in ductal carcinoma in situ of the breast. *J Natl Cancer Inst Monogr* 2010; 2010: 162–177.
- Narod SA, Iqbal J, Giannakeas V et al. Breast cancer mortality after a diagnosis of ductal carcinoma in situ. *JAMA Oncol* 2015; 1: 888–896.
- Lebeau A, Kuhn T. Updates in the treatment of ductal carcinoma in situ of the breast. *Curr Opin Obstet Gynecol* 2016; 28: 49–58.
- Francis A, Fallowfield L, Rea D. The LORIS Trial: addressing overtreatment of ductal carcinoma in situ. *Clin Oncol (R Coll Radiol)* 2015; 27: 6–8.
- Elshof LE, Tryfonidis K, Slaets L et al. Feasibility of a prospective, randomised, open-label, international multicentre, phase III, non-inferiority trial to assess the safety of active surveillance for low risk ductal carcinoma in situ — The LORD study. *Eur J Cancer* 2015; 51: 1497–1510.
- Kerlikowske K, Molinaro AM, Gauthier ML et al. Biomarker expression and risk of subsequent tumors after initial ductal carcinoma in situ diagnosis. *J Natl Cancer Inst* 2010; 102: 627–637.

32. Gradishar WJ, Anderson BO, Balssanian R et al. NCCN guidelines insights breast cancer, Version 1.2016. *J Natl Compr Canc Netw* 2015; 13: 1475–1485.
33. Friedrich M, Kuhn T. Ductal carcinoma in situ. In: Diagnosis and treatment of patients with primary and metastatic breast cancer. *AGO Guidelines Breast Version 16.1*. 2016; 401–450. [www.ago-online.de](http://www.ago-online.de).
34. Senkus E, Kyriakides S, Ohno S et al., on behalf of the ESMO Guidelines Committee. Primary breast cancer: ESMO clinical practice guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2015; 26 (Suppl 5): v8–v30.
35. Mitchell KB, Kuerer H. Ductal carcinoma in situ: treatment update and current trends. *Curr Oncol Rep* 2015; 17: 48–53.
36. Van Cleef A, Altintas S, Huizing M et al. Current view on ductal carcinoma in situ and importance of the margin thresholds: A review. *Facts Views Vis Obgyn* 2014; 6: 210–218.
37. Van Zee KJ, Subhedar P, Olcese C et al. Relationship between margin width and recurrence of ductal carcinoma in situ: analysis of 2996 women treated with breast-conserving surgery for 30 years. *Ann Surg* 2015; 262: 623–631.
38. Subhedar P, Olcese C, Patil S et al. Decreasing recurrence rates for ductal carcinoma in situ: analysis of 2996 women treated with breast-conserving surgery over 30 years. *Ann Surg Oncol* 2015; 22: 3273–3281.
39. Shaikh T, Li T, Murphy CT et al. Importance of surgical margin status in ductal carcinoma in situ. *Clin Breast Cancer* 2016; 16: 312–318.
40. Coates AS, Winer EP, Goldhirsch A et al. Tailoring therapies — improving the management of early breast cancer: St Gallen International Expert Consensus on the Primary Therapy of Early Breast Cancer 2015. *Ann Oncol* 2015; 26: 1533–1546.
41. Morrow M, Van Zee KJ, Solin LJ et al. Society of Surgical Oncology–American Society for Radiation Oncology–American Society of Clinical Oncology Consensus Guideline on Margins for Breast-Conserving Surgery With Whole-Breast Irradiation in Ductal Carcinoma in Situ. *J Clin Oncol* 2016; 34: 4040–4046.
42. Qian GW, Ni XJ, Wang Z et al. Effect of radiotherapy on survival of women with locally excised ductal carcinoma in situ of the breast: a Surveillance, Epidemiology, and End Results population-based analysis. *Onco Targets Ther* 2015; 8: 1407–1418.